

UT-Komaba

93-11

July, 1993

Hamiltonian Formulation of the Smooth Bosonization and Local Gauge Symmetry of the Massless Thirring Model

T. Ikehashi

*Institute of Physics, University of Tokyo,
Komaba, Meguro-Ku, Tokyo 153, Japan*

Abstract: Using the hamiltonian formalism, we investigate the smooth bosonization method in which bosonization and fermionization are carried out through a specific gauge-fixing of an enlarged gauge invariant theory. The generator of the local gauge symmetry, which cannot be derived from the lagrangian of the enlarged theory, is obtained by making a canonical transformation. We also show that the massless Thirring model possesses a similar local gauge symmetry for a specific value of the coupling constant.

A few years ago, Alfaro and Damgaard [1] developed an idea that field-enlarging transformations can be used to find equivalent formulations of the same theory. Recently, Damgaard, Nielsen and Sollacher [2] have applied this idea to the 1+1 dimensional abelian bosonization. Their procedure consists of two steps: First, they introduced a bosonic field to a fermionic theory through a chiral transformation. The theory thus enlarged possesses a local gauge symmetry. Second, they showed that a particular gauge-fixing condition enables us to interpolate smoothly between fermionic and bosonic formulations. The accomplishment of this program seems to rely deeply on the use of the path-integral method since the chiral Jacobian of the fermionic measure plays an important role in forming the local gauge symmetry. Another reason which makes one to prefer the path-integral method is that the generator of the local gauge symmetry, which is crucial in the hamiltonian formalism, cannot be found from the enlarged lagrangian. In this letter, we shall show that it is nevertheless possible to formulate the scheme of Damgaard et. al. in the hamiltonian formalism.

First, we would like to clarify the correspondent of the Jacobian in the hamiltonian formulation. We start from a quantized free Dirac fermion with the lagrangian density[†]

$$\mathcal{L} = i\bar{\Psi}\gamma^\mu\partial_\mu\Psi. \quad (1)$$

Let us suppose that the axial vector current is so regularized that under a local chiral transformation $\Psi(x) \rightarrow e^{-ig\gamma_5\phi(x)}\Psi(x)$, $\bar{\Psi}(x) \rightarrow \bar{\Psi}(x)e^{-ig\gamma_5\phi(x)}$ (g being a coupling constant), it transforms as

$$J_5^\mu = \bar{\Psi}\gamma^\mu\gamma_5\Psi \rightarrow J_5^\mu + \frac{g\lambda}{\pi}\partial^\mu\phi, \quad (2)$$

where a parameter λ represents an ambiguity of the regularization [3, 4]. Now let us apply the chiral transformation to the lagrangian by means of iterating infinitesimal chiral rotations. Then, the first infinitesimal transformation yields

$$\mathcal{L} \rightarrow i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + g\bar{\Psi}\gamma^\mu\gamma_5\Psi\partial_\mu(\delta\phi). \quad (3)$$

Noting that the axial vector current in (3) is also transformed by further infinitesimal transformations, we obtain the transformed lagrangian

$$\mathcal{L} = i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + g\bar{\Psi}\gamma^\mu\gamma_5\Psi\partial_\mu\phi + \frac{1}{2}\frac{g^2\lambda}{\pi}\partial_\mu\phi\partial^\mu\phi. \quad (4)$$

The present procedure just parallels what Roskies and Schaposnik [5] showed in the path-integral method. The last term of (4) corresponds to the contribution of the Jacobian. This shows that, in the hamiltonian formalism, the change of the regularized current represents the effect of the Jacobian. The explicit form of the regularization will be presented soon.

In the path-integral formulation, the function $\phi(x)$ can be promoted to a dynamical variable by integrating over ϕ . As a result, the enlarged lagrangian (4) exhibits a local gauge symmetry

$$\phi(x) \rightarrow \phi(x) - \alpha(x), \quad \Psi(x) \rightarrow e^{-ig\gamma_5\alpha(x)}\Psi(x), \quad \bar{\Psi}(x) \rightarrow \bar{\Psi}(x)e^{-ig\gamma_5\alpha(x)}, \quad (5)$$

$$J_5^\mu(x) \rightarrow J_5^\mu(x) + \frac{g\lambda}{\pi}\partial^\mu\alpha(x). \quad (6)$$

[†]Our conventions are $g_{\mu\nu} = \text{diag}(1, -1)$, $\gamma_5 = \gamma^0\gamma^1$, $\gamma^\mu\gamma_5 = \epsilon^{\mu\nu}\gamma_\nu$, $\epsilon_{01} = -\epsilon^{01} = 1$.

In the hamiltonian formulation, however, it is not clear how to promote ϕ to be dynamical at this stage since Ψ is *already* quantized and is a free Dirac fermion. To circumvent this difficulty, we decide to start from a classical theory defined by a lagrangian density

$$\mathcal{L} = i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + g\bar{\Psi}\gamma^\mu\gamma_5\Psi\partial_\mu\phi + \frac{1}{2}\partial_\mu\phi\partial^\mu\phi. \quad (7)$$

By comparison with (4), it is expected that the quantized system exhibits a local gauge symmetry (5) and (6) when currents are regularized to satisfy (6) and when $g^2\lambda/\pi = 1$. The lagrangian density (7) with the mass term $\frac{1}{2}m^2\phi^2$ is known as the RS model [4, 6] and is solved for $g^2\lambda/\pi < 1$. (We ignore the infrared divergence of the massless free boson since we are interested in boson-fermion correspondence at the operator level.) The classical equation of motion has a solution $\Psi(x) = e^{ig\gamma_5\phi(x)}\psi(x)$, where $\psi(x)$ is a free fermion ($i\gamma^\mu\partial_\mu\psi = 0$) and $\phi(x)$ is a harmonic field ($\square\phi = 0$). The canonical momenta conjugate to ϕ is given by

$$\Pi = g\bar{\Psi}\gamma^0\gamma_5\Psi + \partial^0\phi. \quad (8)$$

Quantization is performed by imposing the equal-time canonical commutation relations;

$$\begin{aligned} [\Pi(x), \phi(y)] &= -i\delta(x^1 - y^1), \quad \{\Psi_\alpha(x), \bar{\Psi}_\beta(y)\} = (\gamma_0)_{\alpha\beta}\delta(x^1 - y^1), \\ [\phi(x), \phi(y)] &= [\phi(x), \Psi(y)] = \{\Psi_\alpha(x), \Psi_\beta(y)\} = 0. \end{aligned} \quad (9)$$

We define the regularization of current as [4]

$$J^\mu(x) = \lim_{\varepsilon \rightarrow 0} \bar{\Psi}(x + \varepsilon)\gamma^\mu \exp \left[ig \int_x^{x+\varepsilon} dy_\rho \{ \lambda \epsilon^{\rho\nu} \partial_\nu \phi + (\lambda + 1) \gamma_5 \partial^\rho \phi \} \right] \Psi(x) - \text{v.e.v.}, \quad (10)$$

where v.e.v. stands for the “vacuum expectation value”. The vector and axial vector currents thus regularized can be rewritten in terms of free fermion currents $j^\mu =: \bar{\psi}\gamma^\mu\psi :$, $j_5^\mu =: \bar{\psi}\gamma^\mu\gamma_5\psi :$ and free boson ϕ by inserting the normal ordered solution;

$$J^\mu = j^\mu - \frac{g\lambda}{\pi} \epsilon^{\mu\nu} \partial_\nu \phi, \quad (11)$$

$$J_5^\mu = j_5^\mu - \frac{g\lambda}{\pi} \partial^\mu \phi. \quad (12)$$

Using (12), one can show that the classical equation of motion for ϕ is modified to $(1 - g^2\lambda/\pi)\square\phi = 0$ which disappears at $g^2\lambda/\pi = 1$. It is also worth noting that according to the analysis of the RS model, there is a wavefunction and coupling constant renormalization $\phi = Z^{1/2}\phi^{(\text{ren})}$, $g = Z^{-1/2}g^{(\text{ren})}$ with $Z = (1 - g^2\lambda/\pi)^{-1}$. These results indicate that the model becomes singular at the critical value $g^2\lambda/\pi = 1$. However, the emergence of the local gauge symmetry is not clear from these considerations since there are no first-class constraints (=generator of the local gauge symmetry). To see how the local gauge symmetry appears, it is useful to perform a canonical transformation

$$\Psi(x) =: e^{ig\gamma_5\phi(x)} : \psi(x) \quad (13)$$

inspired by the form of the exact solution. The normal ordering is not needed at $g^2\lambda/\pi = 1$, since the equation of motion for ϕ vanishes at this value. By this transformation, the lagrangian density is rewritten as

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi + \frac{1}{2}\left(1 - \frac{g^2\lambda}{\pi}\right)\partial_\mu\phi\partial^\mu\phi. \quad (14)$$

Note that insertion of (13) into the lagrangian should be done *gradually* as in the derivation of (4). The canonical momenta of ϕ defined from this lagrangian density is

$$\pi^\phi = (1 - \frac{g^2\lambda}{\pi})\partial^0\phi. \quad (15)$$

The canonical variables are now $(\bar{\psi}, \psi, \pi^\phi, \phi)$ with the canonical commutation relations;

$$\begin{aligned} [\pi^\phi(x), \phi(y)] &= -i\delta(x^1 - y^1), \quad \{\psi_\alpha(x), \bar{\psi}_\beta(y)\} = (\gamma_0)_{\alpha\beta}\delta(x^1 - y^1), \\ [\phi(x), \phi(y)] &= [\phi(x), \psi(y)] = \{\psi_\alpha(x), \psi_\beta(y)\} = 0. \end{aligned} \quad (16)$$

The definition (15) implies that a first-class constraint $\Phi_1 \equiv \pi^\phi \approx 0$ emerges when $g^2\lambda/\pi = 1$. This is indeed the generator of the local gauge symmetry (5) and (6) as can be verified through (12), (13) and (16).

To understand the situation at $g^2\lambda/\pi = 1$, it may be intuitive to consider a general case. Let us consider a lagrangian density $\mathcal{L}(\phi, \partial_\mu\phi)$. If we introduce a field θ by a canonical transformation

$$\phi = f(\phi', \theta), \quad (17)$$

the Hilbert space gets enlarged and acquires a local gauge symmetry with a first-class constraint [1, 7]

$$\Phi'_1 \equiv \pi' - \Pi' \left(\frac{\partial f}{\partial \phi'} \right)^{-1} \frac{\partial f}{\partial \theta} \approx 0, \quad (18)$$

where $\pi' = \partial\mathcal{L}(f, \partial_\mu f)/\partial\dot{\theta}$ and $\Pi' = \partial\mathcal{L}(f, \partial_\mu f)/\partial\dot{\phi}'$. In the enlarged Hilbert space constructed from $(\Pi', \phi', \pi', \theta)$, the physical state $|\text{phys}'\rangle$ should satisfy $\Phi'_1|\text{phys}'\rangle = 0$ owing to the gauge invariance. On the other hand, if we solve ϕ' from (17) as $\phi' = g(\phi, \theta)$ and consider this as a canonical transformation for the enlarged Hilbert space, we get a first-class constraint

$$\Phi_1 \equiv \pi \approx 0, \quad (19)$$

where $\pi = \partial\mathcal{L}(f, \partial_\mu f)/\partial\dot{\theta}$ with $f = f(g(\phi, \theta), \theta)$. This constraint has appeared simply because our original lagrangian density $\mathcal{L}(\phi, \partial_\mu\phi)$ did not contain θ . The physical state for the transformed Hilbert space (Π, ϕ, π, θ) ($\Pi = \partial\mathcal{L}(\phi, \partial_\mu\phi)/\partial\dot{\phi}$) should satisfy $\Phi_1|\text{phys}\rangle = -i\delta/\delta\theta|\text{phys}\rangle = 0$. This means that only (Π, ϕ) are physical and (π, θ) are redundant degrees of freedom. It is important to recognize here that we can investigate gauge-transformation properties of $(\Pi', \phi', \pi', \theta)$ in terms of Φ_1 , because Π', ϕ' and π' can be solved with respect to Π, ϕ, π and θ . As for the present case, we could not find the first-class constraint of the form (18) but we can nevertheless know the transformation laws for the operators of $(\bar{\Psi}, \Psi, \Pi, \phi)$ using Φ_1 .

We can also examine the gauge-fixing condition of Damgaard et. al. with the help of $\Phi_1 = \pi^\phi \approx 0$. The physical state should satisfy $\Phi_1|\text{phys}\rangle = -i\delta/\delta\phi|\text{phys}\rangle = 0$. In the language of $(\bar{\psi}, \psi, \pi^\phi, \phi)$, it implies that the physical state should be constructed from $(\bar{\psi}, \psi)$. One can translate this into the language of $(\bar{\Psi}, \Psi, \Pi, \phi)$ by the use of (11), (12) and (13). For example, a combination $J_5^\mu + g\lambda/\pi\partial^\mu\phi$ is observable in $(\bar{\Psi}, \Psi, \Pi, \phi)$ since this is equivalent to j_5^μ which belongs to $(\bar{\psi}, \psi)$. Damgaard et. al. used this combination as a guide to gauge-fixing. Their gauge-fixing condition has a form

$$\begin{aligned} \Phi_2(x) &= \int_{x_0}^x d\xi_\mu \left\{ \Delta(J_5^\mu + \frac{g\lambda}{\pi}\partial^\mu\phi) - \frac{g\lambda}{\pi}\partial^\mu\phi \right\} \\ &= \int_{x_0}^x d\xi_\mu \left(\Delta j_5^\mu - \frac{g\lambda}{\pi}\partial^\mu\phi \right) \approx 0, \end{aligned} \quad (20)$$

where Δ is a parameter. The path-independence of this expression is evident if we recall that the axial vector current for free fermion can be written in a form $j_5^\mu = \partial^\mu \varphi$ [8]. Therefore when $\Delta = 1$, we get $\partial_x^\nu \Phi_2 = J_5^\nu(x) \approx 0$ which indicate that the fermionic current cannot be observed. On the other hand, $\Delta = 0$ implies that the bosonic field ϕ vanishes from the physical states. This can be verified in terms of the Dirac brackets [9]. Let us employ a space-like path

$$\Phi_2(x) = \Delta \int_{-\infty}^{x^1} d\xi_1 j_5^1(x^0, \xi^1) - \frac{g\lambda}{\pi} \{\phi(x^0, x^1) - \phi(x^0, -\infty)\} \approx 0. \quad (21)$$

Then, observing that $[\Phi_i(x), \Phi_j(y)] = i(g\lambda/\pi)\epsilon_{ij}\delta(x^1 - y^1)$ ($\epsilon_{12} = -\epsilon_{21} = 1$), we obtain the Dirac bracket for arbitrary operators A and B ,

$$[A(x), B(y)]_D = [A(x), B(y)] + \frac{i\pi}{g\lambda} \int dz^1 [A(x), \Phi_i(z^1)] \epsilon_{ij} [\Phi_j(z^1), B(y)]. \quad (22)$$

Noting that $\Pi = gj_5^0 + \pi\phi$ and using $[j^0(x), j^1(y)] = (i/\pi)\partial_x^1\delta(x^1 - y^1)$, the Dirac brackets for Π, ϕ and fermionic currents lead to

$$[\Pi(x), \phi(y)]_D = -i\frac{\Delta}{\lambda}\delta(x^1 - y^1), \quad (23)$$

$$[J_5^0(x), J_5^1(y)]_D = (1 - \Delta)\frac{i}{\pi}\partial_x^1\delta(x^1 - y^1). \quad (24)$$

The commutators for Π, ϕ and fermionic currents vanish for $\Delta = 0$ and $\Delta = 1$, accordingly. Since the non-existence of the commutators can be interpreted as disappearance of these variables from the physical space, these results support the observation of Damgaard et. al. It is also tempting to calculate the anti-commutation relation for fermionic fields. With the help of

$$[(e^{ig\gamma_5\phi(x)})_{\alpha\beta}, \psi_\gamma(y)]_D = \Delta\frac{i\pi}{\lambda} (\gamma_5 e^{ig\gamma_5\phi(x)})_{\alpha\beta} \psi_\gamma(y) \theta(x^1 - y^1), \quad (25)$$

we find

$$\begin{aligned} \{\Psi_\alpha(x), \bar{\Psi}_\beta(y)\} &= (\gamma_0)_{\alpha\beta} \delta(x^1 - y^1) \\ &+ \Delta\frac{i\pi}{\lambda} \left[\theta(y^1 - x^1) (e^{ig\gamma_5\phi(x)})_{\alpha\gamma} (\bar{\Psi}(y)\gamma_5)_\beta (e^{-ig\gamma_5\phi(x)}\Psi(x))_\gamma \right. \\ &\quad \left. - \theta(x^1 - y^1) (\bar{\Psi}(y)e^{-ig\gamma_5\phi(y)})_\gamma (\gamma_5\Psi(x))_\alpha (e^{ig\gamma_5\phi(y)})_{\gamma\beta} \right]. \end{aligned} \quad (26)$$

The anti-commutation relation reduces to the usual one for $\Delta = 0$, but it does not vanish at $\Delta = 1$ which is contrary to naive expectation. This suggests that while the smooth bosonization method is successful in deriving the bosonization relations for currents (and the mass-term relations which we do not discuss in this letter), the complete elimination of the fermionic field is not possible. In the path-integral method, this may correspond to the fact that one cannot decouple the generating functional into fermionic and bosonic part when there is a source term of a form $\bar{\Psi}\eta + \bar{\eta}\Psi$.

Finally, we would like to point out the similarity between the massless RS model (7) and the massless Thirring model. The lagrangian density of the massless Thirring model

is given by

$$\begin{aligned}\mathcal{L} &= i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + \frac{g}{2}(\bar{\Psi}\gamma^\mu\Psi)^2 \\ &= i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + \bar{\Psi}\gamma^\mu\Psi A_\mu - \frac{1}{2g}A_\mu A^\mu,\end{aligned}\tag{27}$$

where we have introduced an auxiliary field A_μ in the second line. Parametrizing the auxiliary field as $A_\mu = \epsilon_{\mu\nu}\partial^\nu\phi + \partial_\mu\eta$, we get

$$\mathcal{L} = i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + \bar{\Psi}\gamma^\mu\Psi(\epsilon_{\mu\nu}\partial^\nu\phi + \partial_\mu\eta) + \frac{1}{2g}(\partial_\mu\phi\partial^\mu\phi - \partial_\mu\eta\partial^\mu\eta).\tag{28}$$

As can be seen from this expression, this is almost the lagrangian density of the massless RS model except for the existence of the indefinite metric field η and trivial rescaling of g and ϕ . Indeed, as in the case of the massless RS model, the massless Thirring model exhibits a local gauge symmetry for a specific value of the coupling constant g (and a parameter λ representing regularization ambiguity). Instead of repeating arguments in the hamiltonian formalism, let us employ the path-integral method this time. Making a change of variable $\Psi = e^{-i\gamma_5\phi+i\eta}\psi$ in the partition function and taking into account the change of the fermionic measure [4];

$$\mathcal{D}\Psi\mathcal{D}\bar{\Psi} = \mathcal{D}\psi\mathcal{D}\bar{\psi}\exp\left[-\frac{i\lambda}{2\pi}\int d^2x\partial_\mu\phi\partial^\mu\phi\right],\tag{29}$$

we get

$$\begin{aligned}Z &= \int \mathcal{D}\Psi\mathcal{D}\bar{\Psi}\mathcal{D}\phi\mathcal{D}\eta\exp\left[i\int d^2x\left\{i\bar{\Psi}\gamma^\mu\partial_\mu\Psi + \bar{\Psi}\gamma^\mu\Psi(\epsilon_{\mu\nu}\partial^\nu\phi + \partial_\mu\eta)\right.\right. \\ &\quad \left.\left.+ \frac{1}{2g}(\partial_\mu\phi\partial^\mu\phi - \partial_\mu\eta\partial^\mu\eta)\right\}\right] \\ &= \int \mathcal{D}\psi\mathcal{D}\bar{\psi}\mathcal{D}\phi\mathcal{D}\eta\exp\left[i\int d^2x\left\{i\bar{\psi}\gamma^\mu\partial_\mu\psi + \frac{1}{2}\left(\frac{1}{g} - \frac{\lambda}{\pi}\right)\partial_\mu\phi\partial^\mu\phi - \frac{1}{2g}\partial_\mu\eta\partial^\mu\eta\right\}\right].\end{aligned}\tag{30}$$

When $g\lambda/\pi = 1$, the integration over ϕ diverges. This implies the emergence of a local gauge symmetry in the path-integral formulation. The generator of this local gauge symmetry will be $\Phi_1 \equiv \pi^\phi \approx 0$, where π^ϕ is the conjugate momenta of ϕ defined from the lagrangian in the last line of (30). Exploiting the gauge-fixing condition $\Phi_2 \equiv \phi \approx 0$, we may eliminate the redundant degrees of freedom (π^ϕ, ϕ) , and the remaining physical part describes free fermion.

In conclusion, we have showed that a local gauge symmetry which includes the contribution from the fermionic Jacobian can also be treated in the hamiltonian formalism. The generator obtained from the canonical-transformed lagrangian enabled us to use the Dirac method for constrained systems. We have reproduced the results of Damgaard et. al. for bosonization relations of currents. It may be interesting to apply this method to the non-abelian bosonization [10].

Acknowledgements: I would like to thank K. Ohta, K. Itakura, I. Ichinose and H. Mukaida for discussions. I am grateful to K. Ohta for his careful reading of the manuscript.

References

- [1] J. Alfaro and P.H. Damgaard, Ann. Phys. (N.Y.) 202 (1990) 398
- [2] P.H. Damgaard, H.B. Nielsen and R. Sollacher, Nucl. Phys. B385 (1992) 227;
Phys. Lett. B296 (1992) 132 and CERN-preprints CERN-TH-6486/92
- [3] K. Harada, H. Kubota and I. Tsutsui, Phys. Lett. B173 (1986) 77
- [4] R. Banerjee, Z. Phys. C25 (1984) 251
- [5] R. Roskies and F.A. Schaposnik, Phys. Rev. D23 (1981) 558
- [6] K.D. Rothe and I.O. Stamatescu, Ann. Phys. (N.Y.) 95 (1975) 202
- [7] A. Hosoya and K. Kikkawa, Nucl. Phys. B101 (1975) 271
- [8] B. Klaiber, 1967 Boulder Lectures in Theoretical Physics, (Gordon and Breach, New York, 1968);
E. Abdalla, M.C.B. Abdalla and K.D. Rothe, Non-Perturbative Methods in 2 Dimensional Quantum Field Theory (World Scientific, Singapore, 1991) p15
- [9] P.A.M. Dirac, Lectures on Quantum Mechanics, (Belfar Graduate School of Science, Yeshiva Univ., 1964)
- [10] E. Witten, Comm. Math. Phys. 92 (1984) 455